

PRV

PATENT- OCH REGISTRERINGSVERKET
Patentavdelningen

**Intyg
Certificate**

Härmed intygas att bifogade kopior överensstämmer med de handlingar som ursprungligen ingivits till Patent- och registreringsverket i nedannämnda ansökan.

This is to certify that the annexed is a true copy of the documents as originally filed with the Patent- and Registration Office in connection with the following patent application.

(71) Sökande *Biacore AB, Uppsala SE*
Applicant (s)

(21) Patentansökningsnummer *0301058-4*
Patent application number

(86) Ingivningsdatum *2003-04-10*
Date of filing

Stockholm, 2004-02-17

För Patent- och registreringsverket
For the Patent- and Registration Office

Hjördis Segerlund
Hjördis Segerlund

Avgift
Fee 170:-

**CERTIFIED COPY OF
PRIORITY DOCUMENT**

METHOD AND KIT FOR CELL ANALYTE ASSAY

Technical field of the invention

The present invention relates to a method of analysing a cell sample for cell
5 surface and intracellularly bound analytes as well as to a kit for cell characterisation.

Background of the invention

The cell walls of living organisms expose a number of cell surface elements of
different functions such as, for example, receptors that interact with specific signal
10 species in the cell environment, and antigens having specific immune reactions.
Information on the presence and concentration of such surface elements in a cell sample
is sought for various reasons. For instance, many disease states in particular animals,
such as humans, have a genetic basis and can be characterised by changes in the
patterns or levels of expression of various genes. Some of the genetic changes are
15 expressed by alterations in cell surface antigens. For example, leukemia may be
diagnosed on the basis of expression of specific CD, lymphoid and myeloid antigens.

An assay device based on cell surface antigen determination is disclosed in WO
00/39580. The assay device comprises an array of different molecules having a binding
partner in a biological sample, wherein the pattern of binding of the binding partners,
20 which may be cells or components released by the cells, to the respective molecules on
the array provides an indication of a normal condition or disease condition or a
propensity therefor to develop in an animal, avian or plant species, or the presence of a
particular cell type or microbial, viral, parasitic or other pathogenic agent.

WO 96/38729 discloses the use of an inhibition type immunoassay to detect
25 pathogens in a sample suspension. The suspension is first reacted with a predetermined
excess of antibodies against the pathogen, and all pathogens are then removed from the
suspension, e.g. by filtration or centrifugation. The amount of antibody in the resulting
pathogen-free solution is then determined by contacting the solution with a sensor
surface having immobilised receptor to the antibody.

30 WO 90/05305 discloses the use of bifunctional molecules for functionalising
sensing surfaces. Each bifunctional molecule has a function for immobilizing the
molecule to the sensing surface, and a bioselective function for interaction with a target
biomolecule.

It is an object of the present invention to provide an improved array-based method for determining cell surface elements and/or intracellular components, which method does not involve the binding of cells to the array nor require the immobilization of receptor molecules to the array.

5

Summary of the invention

The above and other objects and advantages are obtained by a novel method for analysing a cell sample for cell surface-bound analytes as well as intracellularly bound analytes. The method is based on contacting the cell sample with multiple ligands
10 specific to different receptors on or within the cell, wherein each ligand in addition to the cell-specific part comprises a portion specifically binding to a particular array member. This eliminates the need for purification of cell receptor molecules and the immobilization thereof to sensor surfaces as well as the often problematic regeneration of such receptor-supporting surfaces.

15 In one aspect, the present invention therefore provides a method for analysing a cell sample fluid for cell surface-bound and/or intracellularly bound analytes. This method comprises the steps of:

(i) providing a solid support having on a surface thereof a plurality of different binding agents immobilized at defined positions on the surface, wherein each binding
20 agent comprises one member of a specific binding pair;

(ii) contacting the solid support surface with a set of different ligands, each ligand comprising a first part capable of specifically binding to a specific analyte selected from cell surface-bound analytes and intracellularly bound analytes of a defined cell type, and a second part which comprises the other member of each specific
25 binding pair, such that each ligand binds through its specific binding pair part to a specific position on the solid support surface;

(iii) determining the binding of each ligand to the solid support surface;

(iv) incubating a cell sample-containing fluid with the same set of ligands as in step (ii) to permit the ligands to bind to cell surface-bound or intracellularly bound
30 analytes of cells present in the cell sample fluid;

(v) contacting the cell sample fluid with the solid support surface to permit ligands that have not bound to cell surface-bound or intracellularly bound analytes to bind to the solid support surface;

(vi) determining the binding of each ligand obtained in step (v) and comparing that binding with the binding of the same ligand obtained in step (iii), reduced binding indicating the presence of ligand-specific cell surface-bound analytes or intracellularly bound analytes in the cell sample.

5 In another aspect, the present invention provides an assay kit for cell characterisation, which kit comprises:

(i) a solid support having on a surface thereof an array of different binding agents immobilised at defined positions on the surface, wherein each binding agent comprises one member of a specific binding pair; and

10 (ii) a set of different ligands, each ligand comprising a first part capable of specifically binding to a specific analyte selected from cell surface-bound analytes and intracellularly bound analytes of a defined cell, and a second part which comprises the other member of a respective one of the specific binding pairs, such that each ligand may bind through its specific binding pair part to a specific position on the solid support
15 surface.

In still another aspect, the present invention provides an assay kit for cell characterisation, which kit comprises:

(i) a solid support having on a surface thereof an array of different binding agents immobilized at defined positions on the surface, wherein each binding agent
20 comprises one member of a specific binding pair; and

(ii) a set of binding elements, each of which comprises the other member of a respective one of the specific binding pairs and a reactive group that permits chemical coupling of the binding element to a ligand which is capable of specifically binding to a specific analyte selected from specific cell surface-bound analytes and intracellularly
25 bound analytes of a defined cell.

The above and other aspects of the invention will be evident upon reference to the accompanying drawings and the following detailed description.

Brief description of the drawings

30 Figure 1 is a schematic illustration of a solid support surface having an array of ligand binding agents immobilized thereto.

Figure 2 is the array surface in Fig. 1 with ligands immobilized thereto.

Figure 3 is a schematic illustration of a cell having the same ligands as in Fig. 2 bound to different receptors on the cell surface.

Figure 4 is a diagram which shows detector response versus cell concentration at a sensor surface with immobilized anti-mouse IgG antibody for a fluid containing anti-EGF antibody (at 5 nM concentration) after contact thereof with cells that do not express EGF receptor (Jurkat cells) and cells that express EGF-receptor (A431 cells).

Definitions

In describing and claiming the present invention, the following terminology will be used in accordance with the definitions set below.

"Array" and "array surface" as used herein are to be interpreted broadly and generally relate to a linear or two-dimensional array of discrete defined regions (here at least two), each having a finite area, formed on a solid support, usually on a continuous surface thereof, and supporting one or more binding agents. Ordered arrays of nucleic acids, proteins, small molecules, cells or other substances on a solid support enable parallel analysis of complex biochemical samples. In a "microarray", the density of discrete regions, or spots, is typically at least 100/cm², and the discrete regions typically have a diameter in the range of about 10-500 μ m and are separated from other regions in the array by about the same distance.

"Defined region" as used herein relates to a localized area on the solid support surface. The defined region may have any desired shape, such as circular, rectangular, elliptical, etc, and is often referred to as a "spot".

"Solid support" as used herein is meant to comprise any solid (flexible or rigid) substrate onto which it is desired to apply an array of one or more binding agents. The substrate may be biological, non-biological, organic, inorganic or a combination thereof, and may be in the form of particles, strands, precipitates, gels, sheets, tubings, spheres, containers, capillaries, pads, slices, films, plates, slides, etc, having any convenient shape, including disc, sphere, circle, etc. The substrate surface supporting the array may have any two-dimensional configuration and may include, for example steps, ridges, kinks, terraces and the like and may be the surface of a layer of material different from that of the rest of the substrate.

"Specific binding pair" (abbreviated "sbp") as used herein describes a pair of molecules (each being a member of a specific binding pair) which are naturally derived

or synthetically produced. One of the pair of molecules has a structure (such as an area or cavity) on its surface that specifically binds to (and is therefore defined as complementary with) a particular structure (such as a spatial and polar organisation) of the other molecule, so that the pair have the property of binding specifically to each other. Examples of types of specific binding pairs (without any limitation thereto) are antigen-antibody, antibody-hapten, biotin-avidin, ligand-receptor (e.g. hormone receptor, peptide-receptor, enzyme-receptor), carbohydrate-protein, carbohydrate-lipid, lectin-carbohydrate, nucleic acid-nucleic acid (such as oligonucleotide-oligonucleotide).

“Nucleic acid” refers to a deoxyribonucleotide polymer (DNA) or ribonucleotide polymer (RNA) in either single- or double-stranded form, and also encompasses synthetically produced analogs that can function in a similar manner as naturally occurring nucleic acids. While natural nucleic acids have a phosphate backbone, artificial nucleic acids may contain other types of backbones, nucleotides or bases. These include, for instance, peptide nucleic acids (PNAs) as described in e.g. US-A-5,948,902 and the references cited therein; pyranosyl nucleic acids (p-NAs) as described in e.g. WO 99/15540 (p-RNAs), WO 99/15539 (p-RNAs), and WO 00/11011 (p-DNAs); locked nucleic acids (LNAs), as described in e.g. US-A-6,316,198; and phosphothionates and other variants of the phosphate backbone of native nucleic acids.

“Oligonucleotide” refers to single stranded nucleotide multimers of from about 5 to about 100 nucleotides.

“Ligand” as used herein refers to a heterobifunctional molecule that, on one hand, has affinity for a given cell surface-bound or intracellularly bound analyte and, on the other hand, is capable of specifically binding to a moiety immobilized on a solid support. The ligand may be a naturally occurring molecule or one that has been synthesized.

“Antibody” as used herein means an immunoglobulin which may be natural or partly or wholly synthetically produced and also includes active fragments, including Fab antigen-binding fragments, univalent fragments and bivalent fragments. The term also covers any protein having a binding domain which is homologous to an immunoglobulin binding domain. Such proteins can be derived from natural sources, or partly or wholly synthetically produced. Exemplary antibodies are the immunoglobulin isotypes and the Fab, Fab', F(ab')₂, scFv, Fv, dAb, and Fd fragments.

"Hapten" as used herein means a low molecular species that may give rise to an immune response only when coupled to a larger molecule or cell or by aggregation. After immunisation, however, free haptens may react with antibodies.

5 "Cell surface analyte" as used herein refers to a molecule situated on the external surface of a cell. The cell surface analyte may be an antigen having a specific immune reaction. Cell surface antigens may, for example, consist of carbohydrates, lipids or proteins.

10 "Intracellularly bound analyte" as used herein refers to a molecule situated inside a cell and bound to an intracellular structure in such a way that it can not freely diffuse out of a cell wall that is permeable to specific binding partners. The intracellularly bound analyte may be an antigen having a specific immune reaction. Intracellularly bound analytes may, for example, consist of carbohydrates, lipids or proteins.

15 "Cell sample" as used herein means any sample of cells either from cell culture or isolated from an organism, an organ, a body liquid or a tissue. A cell sample preparation may be in the form of a suspension or adherent to a surface. The cell sample preparation may contain intact cells, or cells where the cell membrane has been rendered permeable so that specific binding partners to intracellular components can enter and leave the cell interior. This situation can for instance be achieved by fixation and permeabilization methods as described by Hedley, D.W., et al. in Cytometry, 20 (Communications in clinical cytometry) 46:72-78 (2001).

The terms "set" and "plurality" as used herein, means at least two.

25 In the specification and the appended claims, the singular forms "a", "an", and "the" are meant to include plural reference unless it is stated otherwise. Also, unless defined otherwise, technical and scientific terms used herein have the same meaning as commonly understood to a person skilled in the related to the invention.

Detailed description of the invention

30 As mentioned above, the present invention relates to a method of analysing a cell sample for a set of cell surface-bound analytes or intracellularly bound analytes as well as to an array surface that can be used therefor. Basically, the invention resides in (i) providing an array of immobilised specific binding agents for a set of different ligands, where each ligand is specific to a respective cell surface-bound or

intracellularly bound analyte, and (ii) using the array and the set of ligands in an inhibition type assay format to determine cell surface-bound or intracellularly bound analytes in the cell sample. While the following description is primarily written with respect to cell surface-bound analytes, it is understood that the description is equally applicable to intracellularly bound analytes.

An array surface for use in the invention may be constructed by immobilising to defined areas of the surface a number of different ligand recognition elements, which for the sake of illustration are named A, B, C, D..., and each of which is one member of a specific binding pair. A schematic illustration of an array surface with immobilized recognition elements A, B, C, D... is shown in Fig. 1. The corresponding other members of these specific binding pairs, here named a, b, c, d..., are each attached (such as conjugated) to a respective one of a set of different ligands L1, L2, L3, L4..., which are directed against specific surface structures or elements (such as e.g. antigens) that may be present on the surface of a particular cell to form modified ligands aL1, bL2, cL3, dL4... The specific structures or elements on the cell surface are below often referred to as analytes. When the modified ligands are contacted with the array surface, aL1 combines with A, bL2 combines with B, etc, as is schematically illustrated in Fig. 2. The specific binding of a modified ligand to its recognition element on the array surface may be detected by a variety of different methods well known in the art as will be described in more detail below.

To carry out a study or "mapping" of a cell surface, the array surface is first contacted with the ligands, each ligand preferably at a certain known concentration, and the responses at the different recognition elements A, B, C, D... are measured. Then, preferably the same concentrations of ligands are incubated with a desired cell sample, which, for example, may be intact cells, expressing surface-bound analytes for the ligands aL1, bL2 etc. During the incubation, reactions will occur in solution and the ligands will bind to their respective cell surface analytes, as is schematically illustrated in Fig. 3. The sample solution is then brought in contact with the array surface which, as described above, permits self-orientation of the ligands.

Preferably, however, prior to contacting the sample solution with the array surface, suspended or emulsified matter including cells, membranes etc are removed from the sample solution by, for instance, filtration or centrifugation. Alternatively, the sample solution may be passed over the array surface in a laminar flow that

concentrates the suspended or emulsified matter in a central flow core spaced from the surface as described in WO 95/27208 (the disclosure of which is incorporated by reference herein). These procedures aim at separating free ligands aL1, bL2 etc from ligand(s) bound to cell surface specific analyte(s).

5 After the incubation of the cell sample with the ligand set, at least some of the different ligands will, depending on the presence and amount of corresponding cell surface analytes, be bound to the respective cell surface analytes, and optionally be removed along with the cells in the separation step. When the (optionally separated) cell sample solution is contacted with the array surface, the binding of the ligands aL1, bL2
10 etc to the corresponding recognition elements A, B etc on the array surface, will differ in rate and extent in comparison with the first-mentioned situation where the ligands were contacted with the array surface without preceding exposure to their respective cells surface analytes. Thus, when a difference in the binding of a specific ligand to the array surface is observed, this is an indication that the corresponding analyte was
15 present in the cell sample. In this way at least a qualitative determination of cell surface analytes may be obtained.

As is readily recognised, repeating the above procedure with different concentrations of the cells or membranes may provide a quantitative measure of cell surface analytes in a cell sample, such as e.g. the number of a particular receptor per
20 cell.

The procedure described above may thus be used to simultaneously determine or monitor a large number of cell surface-bound as well as intracellularly bound analytes and their concentrations.

I many cases, it may be advantageous to carry out the assay procedure in one or
25 more flow cells (arranged in series or in parallel). Flow cell designs that may be contemplated are well known to the skilled person and need not be described herein. It is to be noted that an array may be distributed between two or more flow cells. For example, an array of eight ligands may be provided in four flow cells, i.e. two ligands in each flow cell.

30 Specific binding pairs Aa, Bb, Cc, Dd... that may be used are well known to the skilled person and may, for example, be hybridising strands of nucleic acids, usually oligonucleotides, one of the strands being linked (such as conjugated) to the ligands L1, L2, L3, L4.... As another example may be mentioned antibodies or antibody fragments

A, B, C, D... directed against haptens a, b, c, d... that are linked (such as conjugated) to the ligands L1, L2, L3, L4....Such heterobifunctional ligands and the preparation thereof are *per se* well known to the skilled person and need not be described any further herein.

5 A kit for carrying out the above described assay procedure may comprise a sensor surface, such as a sensor chip, with immobilized recognition elements A, B, C, D..., and a set of corresponding cell surface antigen specific ligands aL1, bL2, cL3, dL4....(or intracellularly bound antigen specific ligands).

10 Alternatively, the kit may comprise a sensor surface, such as a sensor chip, with immobilized recognition elements A, B, C, D..., and a set of corresponding binding elements a,b,c,d...that have been modified with a reactive group (e.g. succinimide ester, maleimide, dipyridyl-disulfide) so that they can readily be incorporated into ligands L1, L2 ,L3, L4...of choice.

15 Exemplary cell types that may be of interest for use in the assay include: liver cells, gastrointestinal cells, epithelial cells, endothelial cells, kidney cells, cancer cells, blood cells, stem cells, bone cells, smooth muscle cells, striated muscle cells, cardiac muscle cells, nerve cells. Blood cells include e.g. leukocytes, such as neutrophils, lymphocytes, monocytes, eosiniphils, basophils, macrophages.

20 Methods for detecting the presence of bound ligands on the array surface may be chosen from a wide variety of detection techniques, including, for example, marker-based techniques, where the analyte(s) or an analyte specific reagent is labelled, e.g. with a radiolabel, a chromophore, fluorophore, chemiluminescent moiety or a transition metal.

25 For many applications, detection may be performed with a biosensor. A biosensor is broadly defined as a device that uses a component for molecular recognition (for example a layer with immobilised antibodies) in conjunction with a solid state physicochemical transducer. Biosensors may be based on a variety of detection methods. Typically such methods include, but are not limited to, mass detection methods, such as piezoelectric, optical, thermo-optical and surface acoustic
30 wave (SAW) device methods, and electrochemical methods, such as potentiometric, conductometric, amperometric and capacitance methods. With regard to optical detection methods, representative methods include those that detect mass surface concentration, such as reflection-optical methods, including both internal and external

reflection methods, angle, wavelength or phase resolved, for example ellipsometry and evanescent wave spectroscopy (EWS), the latter including surface plasmon resonance (SPR) spectroscopy, Brewster angle refractometry, critical angle refractometry, frustrated total reflection (FTR), evanescent wave ellipsometry, scattered total internal reflection (STIR), optical wave guide sensors, evanescent wave-based imaging such as critical angle resolved imaging, Brewster angle resolved imaging, SPR angle resolved imaging, and the like. Further, photometric methods based on, for example, evanescent fluorescence (TIRF) and phosphorescence may also be employed, as well as waveguide interferometers.

One exemplary type of SPR-based biosensors is sold by Biacore AB (Uppsala, Sweden) under the trade name BIACORE® and is used in the Example below. These biosensors utilize a SPR based mass-sensing technique to provide a "real-time" binding interaction analysis between a surface bound ligand and an analyte of interest.

A detailed discussion of the technical aspects of the BIACORE® instruments and the phenomenon of SPR may be found in U.S. Patent No. 5,313,264. More detailed information on matrix coatings for biosensor sensing surfaces is given in, for example, U.S. Patents Nos. 5,242,828 and 5,436,161. In addition, a detailed discussion of the technical aspects of the biosensor chips used in connection with the BIACORE® instruments may be found in U.S. Patent No. 5,492,840. The full disclosures of the above-mentioned U.S. patents are incorporated by reference herein.

In the following Example, various aspects of the present invention are disclosed more specifically for purposes of illustration and not limitation.

Example

Instrumentation

A BIACORE® 3000 instrument (Biacore AB, Uppsala, Sweden) was used. BIACORE® instruments are based on surface plasmon resonance (SPR) detection at gold surfaces, and a micro-fluidic system is used for passing samples and running buffer through four individually detected flow cells (one by one or in series), with very high precision and with small sample volumes needed. As sensor chip was used Series S CM5 (Biacore AB, Uppsala, Sweden) which has a gold-coated surface with a covalently linked carboxymethyl-modified dextran polymer hydrogel. The output from the instrument is a "sensorgram" which is a plot of detector response (measured in

“resonance units”, RU) as a function of time. An increase of 1000 RU corresponds to an increase of mass on the sensor surface of approximately 1 ng/mm².

Preparation of cell samples

5 A 431 human epidermoid carcinoma cell line (ATCC #CRL-2592) which overexpresses epidermal growth factor (EGF), and a human T-cell leukaemia Jurkat cell line (ATCC#CRL-10915) which does not express the EGF receptor were each grown in culture media. A431 cells were grown in DMEM/FCS and Jurkat cells in RPMI/Lglu/FCS. Jurkat cells were harvested after three days, and A431 cells when they
10 were 90-100% confluent. A431 cells were detached using Versene (EDTA solution) at a 1:5000 dilution.

After washing with PBS (10 mM phosphate buffer pH 7.4 with 150 mM NaCl) and PBS containing 2% bovine serum albumin (BSA, Sigma A3059), cells were serially diluted into microtiter well plates.

15

Assay for cell surface EGF

Mouse anti-EGFr antibody (obtained from Oncogene) at a specific concentration of 5 nM, was added to each cell sample well and incubated for 30 minutes on a rocking plate. The plate was then centrifuged for three minutes at 1400 rpm. The supernatants
20 were injected into the BIACORE® 3000 over sensor chip surfaces with rabbit anti-mouse IgG1 antibody (obtained from Biacore AB, BR-1000-55) immobilized (18400 RU) to detect free antibody in the supernatants, i.e. antibody that had not bound to cell surface EGF. Cell samples were injected for two minutes. Between cell sample injections the sensor surface was regenerated by an injection of 10 mM glycine-HCl at
25 pH 1.2 for 10 seconds.

The results are shown in Fig. 4. As can be seen from the graph, the level of captured antibody when Jurkat cells were used was constant and independent of cell concentration. In contrast the level of captured antibody was reduced when A431 cells were used and more so at higher cell concentration, reflecting that the antibody was
30 partially bound to EGF-receptor on cell surfaces.

Claims

1. A method for analysing a cell sample for cell surface-bound or intracellularly bound analytes, which method comprises the steps of:
 - 5 (i) providing a solid support having on a surface thereof a plurality of different binding agents immobilized at defined positions on the surface, wherein each binding agent comprises one member of a specific binding pair;
 - (ii) 10 contacting the solid support surface with a set of different ligands, each ligand comprising a first part capable of specifically binding to a specific analyte selected from cell surface-bound analytes and intracellularly bound analytes of a defined cell type, and a second part which comprises the other member of each specific binding pair, such that each ligand binds through its specific binding pair part to a specific position on the 15 solid support surface;
 - (iii) determining the binding of each ligand to the solid support surface;
 - (iv) incubating a cell sample-containing fluid with the same set of ligands as in step (ii) to permit the ligands to bind to cell surface-bound or intracellularly bound analytes of cells present in the cell sample fluid;
 - 20 (v) contacting the cell sample fluid with the solid support surface to permit ligands that have not bound to cell surface-bound or intracellularly bound analytes to bind to the solid support surface;
 - (vi) 25 determining the binding of each ligand obtained in step (v) and comparing that binding with the binding of the same ligand obtained in step (iii), reduced binding indicating the presence of ligand-specific cell surface-bound analytes or intracellularly bound analytes in the cell sample.
2. 30 The method according to claim 1, wherein cells and cell fragments are removed from the cell sample before contacting the incubated cell sample fluid with the solid support surface.

3. The method according to claim 2, wherein the cells and fragments are removed by filtration or centrifugation.
4. The method according to claim 1, 2 or 3, wherein each ligand in step (ii) of claim 1 is used at a known concentration, and that the same concentration of each ligand is used in step (v) of claim 1.
5. The method according to claim 4, wherein steps (v) and (vi) of claim 1 are repeated with at least one different cell concentration of the cell sample-containing fluid, and that quantitative measures of cell surface-bound analytes or intracellularly bound analytes are determined.
6. The method according to any one of claims 1 to 5, wherein the cell sample contains intact cells.
7. The method according to any one of claims 1 to 5, wherein the cell sample contains cells, the membranes of which have been made permeable to ligands.
8. The method according to any one of claims 1 to 7, wherein binding to the solid support surface is detected by a label-free detection method.
9. The method according to claim 8, wherein the detection method is based on mass-sensing, especially based on evanescent wave sensing, for example surface plasmon resonance (SPR).
10. The method according to any one of claims 1 to 9, wherein the solid support surface is provided in at least one flow cell.
11. An assay kit for cell characterization, comprising:
 - (i) a solid support having on a surface thereof an array of different binding agents immobilized at defined positions on the surface, wherein each binding agent comprises one member (A, B, C, D...) of a specific binding pair (aA, bB, cC, dD...); and

- (ii) a set of different ligands (aL1, bL2, cL3, dL4...), each ligand comprising a first part (L1, L2, L3, L4...) capable of specifically binding to an analyte selected from specific cell surface-bound analytes and intracellularly bound analytes of a defined cell, and a second part which comprises the other member (a, b, c, d...) of a respective one of the specific binding pairs, such that each ligand may bind through its specific binding pair part to a specific position on the solid support surface.
- 10 12. The assay kit according to claim 11, wherein the specific binding pairs comprise complementary nucleic acid strands, especially oligonucleotides.
13. The assay kit according to claim 11, wherein the specific binding pairs comprise hapten/antibody.
- 15 14. The assay kit according to claim 13, wherein the specific binding pair part of the ligand comprises a hapten.
15. The assay kit according to any one of claims 11 to 14, wherein the solid support surface is a sensor surface, preferably a sensing surface of a biosensor.
- 20 16. An assay kit for cell characterization, comprising:
- 25 (i) a solid support having on a surface thereof an array of different binding agents immobilized at defined positions on the surface, wherein each binding agent comprises one member (A, B, C, D...) of a specific binding pair (aA, bB, cC, dD...); and
- 30 (ii) a set of binding elements, each of which comprises the other member (a, b, c, d...) of a respective one of the specific binding pairs (aA, bB, cC, dD...) and a reactive group that permits chemical coupling of the binding element to a ligand (L1, L2, L3, L4...) which is capable of specifically binding to a specific analyte selected from specific cell surface-bound analytes and intracellularly bound analytes of a defined cell.

- 10

Abstract

The present invention relates to a method for analysing a cell sample for cell surface-bound or intracellularly bound analytes by providing an array of immobilised specific binding agents for a set of different ligands, where each ligand is specific to a respective cell surface-bound or intracellularly bound analyte, and using the array and the set of ligands in an inhibition type assay format to determine cell surface-bound or intracellularly bound analytes in the cell sample. The invention also relates to assay kits for cell characterization.

10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
220




FIG. 1




FIG. 2





FIG. 3

**FIG. 4**